

# Some graded analogues of one-parameter subgroups and applications to the cohomology of $GL_{m|n}(r)$

---

Christopher M. Drupieski, DePaul University

Joint work with Jonathan Kujawa, University of Oklahoma

AMS Fall Southeastern Section Meeting, 23 September 2017

Special Session on Categorical Methods in Representation Theory

Work over a field  $k$  of characteristic  $p > 0$ . (Assume  $k = \bar{k}$  and  $p \geq 3$ .)

$\mathbb{G}_a$  – additive group scheme

$$\mathbb{G}_a(A) = (A, +)$$

$\mathbb{G}_{a(r)}$  –  $r$ -th Frobenius kernel of  $\mathbb{G}_a$

$$\mathbb{G}_{a(r)}(A) = \{t \in A : t^{p^r} = 0\}$$

### Infinitesimal one-parameter subgroups

Given an affine group scheme  $G$ , an infinitesimal one-parameter subgroup of height  $\leq r$  in  $G$  is a group scheme homomorphism

$$\nu : \mathbb{G}_{a(r)} \rightarrow G.$$

The set of all such homomorphisms is denoted  $V_r(G)$ .

$$V_r(G) = \mathbf{Hom}_{\text{Grp}}(\mathbb{G}_{a(r)}, G)$$

The set  $V_r(G)$  admits the structure of an affine variety.

Let  $A$  be an augmented  $k$ -algebra. Suppose  $H^\bullet(A, k) = \text{Ext}_A^\bullet(k, k)$  is “commutative” and finitely generated.

### Cohomological spectrum and support varieties

The **cohomological spectrum** of  $A$  is the affine algebraic variety

$$|A| = \text{MaxSpec} \left( H^\bullet(A, k) \right).$$

Given an  $A$ -module  $M$ , let  $I_A(M)$  be the kernel of the map

$$H^\bullet(A, k) = \text{Ext}_A^\bullet(k, k) \xrightarrow{-\otimes M} \text{Ext}_A^\bullet(M, M).$$

The **cohomological support variety** associated to  $M$  is

$$|A|_M = \text{MaxSpec} \left( H^\bullet(A, k) / I_A(M) \right),$$

a closed subvariety of the cohomological spectrum.

In reasonable settings, support varieties detect projectivity.

### Suslin–Friedlander–Bendel (1997)

Let  $G$  be an infinitesimal group scheme of height  $\leq r$ , and let  $kG$  be its group ring ( $kG = k[G]^\#$ ). Then there exists a homeomorphism

$$|kG| \cong V_r(G) = \mathbf{Hom}_{\mathit{Grp}}(\mathbb{G}_{a(r)}, G)$$

For  $G = GL_{n(r)}$ , the  $r$ -th Frobenius kernel of  $GL_n$ , one has

$$V_r(GL_{n(r)}) = \{(\alpha_0, \dots, \alpha_{r-1}) \in \mathfrak{gl}_n^{\times r} : \alpha_i^p = 0, [\alpha_i, \alpha_j] = 0, \forall i, j\}.$$

If  $G$  is an arbitrary infinitesimal group scheme of height  $\leq r$ , then there exists a closed embedding  $G \hookrightarrow GL_{n(r)}$  for some  $n$ .

If  $\nu : \mathbb{G}_{a(r)} \rightarrow G$  is a one-parameter subgroup, and if  $M$  is a rational  $G$ -module, then  $M$  pulls back to a rational  $\mathbb{G}_{a(r)}$ -module,  $\nu^*(M)$ .

Equivalently,  $\nu^*(M)$  is a module over the group algebra

$$k\mathbb{G}_{a(r)} = k[\mathbb{G}_{a(r)}]^\# = k[u_0, \dots, u_{r-1}]/(u_0^p, \dots, u_{r-1}^p).$$

### Suslin–Friedlander–Bendel (1997)

Let  $G$  be infinitesimal of height  $\leq r$ . If  $M$  is a finite-dimensional rational  $G$ -module, then

$$|kG|_M \cong \{ \nu \in V_r(G) : \nu^*(M) \text{ is not free over } k[u_{r-1}]/(u_{r-1}^p) \}.$$

So, for example, the projectivity of  $M$  can be detected by restrictions along various  $\nu$  to algebras of the form  $k[u]/(u^p)$ .

## Our motivating question

(How) can this be generalized to **supergroups**?

## Wikipedia definition of a supergroup

A **supergroup** is a music group whose members are already successful as solo artists or as part of other groups or well known in other musical professions.

## What do we mean by “super”?

An object is “super” if it is appropriately graded by  $\mathbb{Z}/2\mathbb{Z} = \{\bar{0}, \bar{1}\}$ .

- Super vector spaces  $V = V_{\bar{0}} \oplus V_{\bar{1}}$
- $V \otimes W \cong W \otimes V$  via the **supertwist**  $v \otimes w \mapsto (-1)^{\bar{v} \cdot \bar{w}} w \otimes v$

Define (Hopf) superalgebras and ‘super’ (co)commutativity in terms of the “usual diagrams,” but use the supertwist when objects pass.

## Examples of Hopf superalgebras

- Ordinary Hopf algebras (as purely even superalgebras)
- $\mathbb{Z}$ -graded Hopf algebras in the sense of Milnor and Moore
- Enveloping superalgebras of (restricted) Lie superalgebras
- Exterior algebra  $\Lambda(V)$  over a (purely odd) vector space  $V$  (both commutative and cocommutative in the super sense)

## Defining correspondences

affine **super**group schemes  $\overset{\text{anti}}{\longleftrightarrow}$  commutative Hopf **super**algebras

$$G \longleftrightarrow k[G]$$

finite **super**group schemes  $\longleftrightarrow$  f.d. cocommut. Hopf **super**algebras

$$G \longleftrightarrow kG = k[G]^{\#}$$



## Cautionary example

Let  $A = k[u, v]/\langle u^p, v^2 \rangle$  with  $\bar{u} = \bar{0}$  and  $\bar{v} = \bar{1}$ .

Then  $A$  is a Hopf superalgebra with  $u$  and  $v$  both primitive.

(In fact,  $A$  is the restricted enveloping algebra of a RLSA.)

Define  $M$  to be the  $A$ -supermodule with homogeneous basis

$$\{x_0, \dots, x_{p-1}, y_0, \dots, y_{p-1}\}, \quad x_i \text{ even}, \quad y_i \text{ odd},$$

such that  $u \cdot x_i = x_{i+1}$ ,  $u \cdot y_i = y_{i+1}$ ,  $v \cdot x_i = y_{i+1}$ , and  $v \cdot y_i = x_{i+p-1}$ .

**Claim:**  $M$  is projective over all proper cyclic subalgebras of  $A$ , but is not projective over  $A$  itself. So in contrast to the classical theory, need more than just cyclic subalgebras to detect projectivity.

**What supergroups play the role of  $\mathbb{G}_{a(r)}$  in the super theory?** These will be our graded analogues of one-parameter subgroups.

Let  $f = T^p + \sum_{i=1}^{t-1} a_i T^{p^i} \in k[T]$  be a  $p$ -polynomial (no linear term).

Let  $\eta \in k$  be a scalar.

### The infinitesimal multiparameter supergroup $\mathbb{M}_{r,f,\eta}$

$$k\mathbb{M}_{r,f,\eta} = k[u_0, \dots, u_{r-1}, v] / \langle u_0^p, \dots, u_{r-2}^p, u_{r-1}^p + v^2, f(u_{r-1}) + \eta u_0 \rangle$$

–  $u_0, \dots, u_{r-1}$  are even; coproducts look like they do in  $k\mathbb{G}_{a(r)}$

–  $u_{r-1}^p$  is primitive,  $v$  is an odd primitive generator

For  $r = 1$ , this reduces to

$$k[u, v] / \langle u^p + v^2, f(u) + \eta u \rangle$$

Our result: use these supergroups to generalize the SFB calculation

$$|GL_n(r)| \cong V_r(GL_n(r)) = \mathbf{Hom}_{Grp}(\mathbb{G}_{a(r)}, GL_n(r)).$$

Let  $m, n \in \mathbb{Z}_{\geq 0}$ , and let  $A = A_{\bar{0}} \oplus A_{\bar{1}}$  be a commutative superalgebra.

$\text{Mat}_{m|n}(A)_{\bar{0}}$  consists of all block matrices

$$T = \begin{pmatrix} T_1 & T_2 \\ T_3 & T_4 \end{pmatrix},$$

with  $T_1 \in M_{m \times m}(A_{\bar{0}})$ ,  $T_2 \in M_{m \times n}(A_{\bar{1}})$ ,  $T_3 \in M_{n \times m}(A_{\bar{1}})$ ,  $T_4 \in M_{n \times n}(A_{\bar{0}})$ .

In  $\text{Mat}_{m|n}(A)_{\bar{1}}$ , the parities of the entries are reversed.

### General linear supergroup $GL_{m|n}$ and its Frobenius kernels

For each commutative superalgebra  $A$ ,  $GL_{m|n}(A)$  is the group of all invertible matrices in  $\text{Mat}_{m|n}(A)_{\bar{0}}$ .

$GL_{m|n}(r)$  is the scheme-theoretic kernel of the map that raises each individual matrix entry to the  $p^r$ -th power.

Ambient superscheme (analogue of the commuting variety for  $GL_n$ )

$$V_r(GL_{m|n})(A) = \left\{ (\alpha_0, \dots, \alpha_{r-1}, \beta) \in (\text{Mat}_{m|n}(A)_{\bar{0}})^{\times r} \times \text{Mat}_{m|n}(A)_{\bar{1}} : \right. \\ \left. [\alpha_i, \alpha_j] = [\alpha_i, \beta] = 0 \text{ for all } 0 \leq i, j \leq r-1, \right. \\ \left. \alpha_i^p = 0 \text{ for all } 0 \leq i \leq r-2, \text{ and } \alpha_{r-1}^p + \beta^2 = 0 \right\}.$$

### Lemma

Homomorphisms  $\rho : \mathbb{M}_{r,f,\eta} \otimes_k A \rightarrow GL_{m|n} \otimes_k A$  correspond to points in the closed sub-superscheme

$$V_{r,f,\eta}(GL_{m|n})(A) = \\ \left\{ (\alpha_0, \dots, \alpha_{r-1}, \beta) \in V_r(GL_{m|n})(A) : f(\alpha_{r-1}) + \eta\alpha_0 = 0 \right\}.$$

Note that  $V_r(GL_{m|n})(k) = \bigcup_{f,\eta} V_{r,f,\eta}(GL_{m|n})(k)$ .

## Following SFB to relate $V_r(GL_{m|n})$ to $|GL_{m|n(r)}|$

- Able to construct algebra homomorphisms

$$k[V_r(GL_{m|n})] \xrightarrow{\bar{\phi}} H(GL_{m|n(r)}, k) \xrightarrow{\psi_{r,f,\eta}} k[V_{r,f,\eta}(GL_{m|n})]$$

- Get induced morphisms of varieties

$$\Theta_{r,f,\eta} : V_{r,f,\eta}(GL_{m|n})(k) \xrightarrow{\Psi_{r,f,\eta}} |GL_{m|n(r)}| \xrightarrow{\Phi} V_r(GL_{m|n})(k)$$

- Showed that  $\Theta_{r,f,\eta}$  is the Frobenius morphism composed with the natural inclusion. The  $V_{r,f,\eta}(GL_{m|n})(k)$  cover  $V_r(GL_{m|n})(k)$ , so...

### Theorem (so far, just for $GL_{m|n(r)}$ )

There is a finite surjective morphism of varieties

$$|GL_{m|n(r)}| \rightarrow V_r(GL_{m|n})(k).$$

# Toward a theory of rank varieties

Consider again the algebra  $A := k[u, v]/\langle u^p, v^2 \rangle = k[u]/\langle u^p \rangle \otimes \Lambda(v)$ .

Then  $H^\bullet(A, k) = k[x, y] \otimes \Lambda(\Lambda)$ , hence  $|A| = k^2$ .

Set  $B = k[u, v]/\langle u^p + v^2 \rangle$ . Each point  $(a, \mu) \in |A|$  defines a map

$$\phi : k[u, v]/\langle u^p + v^2 \rangle \rightarrow k[u, v]/\langle u^p, v^2 \rangle$$

such that  $\phi(u) = a \cdot u$  and  $\phi(v) = \mu \cdot v$ .

## Theorem

Let  $V$  be a finite-dimensional  $A$ -supermodule. Then

$$|A|_V = \left\{ (a, \mu) \in |A| : \text{Ext}_B^i(\phi^*V, \phi^*V) \neq 0 \text{ for } \infty\text{-ly many } i \right\}$$