





























































Symmetry and Groups

- ▶ What is symmetry?
- ▶ What is **a** symmetry?
- ▶ How do you *multiply* two symmetries?

Example: Rigid motion symmetries of an equilateral triangle.

<div style="display: flex; align-items: center;"> <div style="writing-mode: vertical-rl; transform: rotate(180deg);">1st</div> <div style="margin-left: 10px;">2nd</div> </div>						
						
						
						
						
						
						

2nd 1st						
						
						
						
						
						
						

Example: Symmetries of a “notched” line.

Example: Addition mod 6.

+	[0]	[1]	[2]	[3]	[4]	[5]
[0]						
[1]						
[2]						
[3]						
[4]						
[5]						

+	[0]	[1]	[2]	[3]	[4]	[5]
[0]	[0]	[1]	[2]	[3]	[4]	[5]
[1]	[1]	[2]	[3]	[4]	[5]	[0]
[2]	[2]	[3]	[4]	[5]	[0]	[1]
[3]	[3]	[4]	[5]	[0]	[1]	[2]
[4]	[4]	[5]	[0]	[1]	[2]	[3]
[5]	[5]	[0]	[1]	[2]	[3]	[4]

Example: Permutations of the set $\{1, 2, 3\}$.

A **group** is a collection of things that can be “multiplied” together. The “multiplication” must satisfy the following properties:

- ▶ For each three things a , b , and c in the collection,


$$a \cdot (b \cdot c) = (a \cdot b) \cdot c$$

- ▶ There is a thing e in the collection so that, for any other thing a in the collection, $e \cdot a = a$ and $a \cdot e = a$.
- ▶ For each thing a in the collection, there is another thing a' in the collection so that $a \cdot a' = e$ and $a' \cdot a = e$.

The greatest mathematical achievement of the 20th century

The greatest mathematical achievement of the 20th century

great (grāt), *adj.* 1. of an extent, amount, or intensity considerably above the normal or average; very large and imposing.



First-generation proof of CFSG:

- ▶ 10,000–15,000 journal pages
- ▶ spread across some 500 separate articles, written by more than 100 mathematicians from around the world
- ▶ most work done between 1950 and early 1980s

Second-generation proof of CFSG (currently underway):

- ▶ shorter, more direct proof of 3,000–4,000 pages
- ▶ volume 1 published in 1994
- ▶ volume 6 published in 2005 (most recent volume)
- ▶ 12 volumes anticipated

- ▶ How are bigger groups built from copies of smaller groups?

2nd 1st						

- ▶ Is \mathbb{Z}_6 made from a group of order 2 and a group of order 3?

The Periodic Table Of Finite Simple Groups

B, C_n, Z_n 1 1		Dynkin Diagrams of Simple Lie Algebras															C_2 2		
$A_n(4), A_n(5)$ A_5 60	$A_1(2)$ $A_1(7)$ 168	A_1 	A_2 	A_3 	A_4 	A_5 	D_n 	E_6 	E_7 	E_8 	F_4 	G_2 	$G_2(2)$ 120	$B_2(3)$ 2160	$C_3(3)$ 108000000	$D_4(2)$ 174182400	${}^2D_4(2^2)$ 197408720	$G_2(2)$ ${}^2A_2(9)$ 6300	C_3 3
$A_n(6), B_n(2^2)$ A_6 360	$G_2(3)$ $A_1(8)$ 336	A_6 	$G_2(3)$ 	A_7 	A_8 	A_9 	A_{10} 	A_{11} 	A_{12} 	A_{13} 	A_{14} 	A_{15} 	$B_2(4)$ 878400	$C_3(5)$ 22080	$D_4(3)$ 416217981600	${}^2D_4(3^2)$ 11131166687328	${}^2A_2(16)$ 62400	C_5 5	
A_7 2520	$A_1(11)$ 660	$E_6(2)$ 21840 879322 609176 276480	$E_7(2)$ 27720 100800 720720 2822400	$E_8(2)$ 33600 120960 907200 3302400	$F_4(2)$ 945 3264 462 16640	$G_2(3)$ 4265486	${}^3D_4(2^3)$ 211341312	${}^2E_6(2^2)$ 76342 479488 79168 194208	${}^2B_2(2^3)$ 29120	${}^2F_4(2)'$ 17971200	${}^2G_2(3^3)$ 10075168 472	$B_3(2)$ 1451520	$C_4(3)$ 68784768 61648000	$D_5(2)$ 214992094800	${}^2D_3(2^2)$ 21619379400 60	${}^2A_2(25)$ 126000	C_7 7		
A_8 20160	$A_1(13)$ 1992	$E_6(3)$ 27720 100800 720720 2822400	$E_7(3)$ 33600 120960 907200 3302400	$E_8(3)$ 40320 141120 1008000 3604800	$F_4(3)$ 1176 3936 576 18432	$G_2(4)$ 251376000	${}^3D_4(3^3)$ 216681166912	${}^2E_6(3^2)$ 32537600	${}^2B_2(2^5)$ 32537600	${}^2F_4(2^3)$ 18616032000 16617945440	${}^2G_2(3^5)$ 49402487 61934082	$B_2(5)$ 6480000	$C_3(7)$ 60419240	$D_4(5)$ 90000000 16148040	${}^2D_4(4^2)$ 47376471	${}^2A_3(9)$ 3265920	C_{11} 11		
A_9 181440	$A_1(17)$ 2448	$E_6(4)$ 33600 120960 907200 3302400	$E_7(4)$ 40320 141120 1008000 3604800	$E_8(4)$ 47040 161280 1120000 3916800	$F_4(4)$ 14112 4536 896 30720	$G_2(5)$ 346908000	${}^3D_4(4^3)$ 67368000 64278640	${}^2E_6(4^2)$ 340933600	${}^2B_2(2^7)$ 2003891624 30234832512	${}^2G_2(3^7)$ 138287680	$B_2(7)$ 9430511482 91130919320	$C_5(9)$ 90000000 16148040	$D_3(3)$ 17400 212160 8000000	${}^2D_4(5^2)$ 5515776	${}^2A_2(64)$ 5515776	C_{13} 13			
A_n at $\frac{n}{2}$	$A_n(q)$ $E_6(q)$ $E_7(q)$ $E_8(q)$ $F_4(q)$ $G_2(q)$ ${}^3D_4(q^3)$ ${}^2E_6(q^2)$ ${}^2B_2(2^{2n+1})$ ${}^2F_4(2^{2n+1})$ ${}^2G_2(3^{2n+1})$	$E_6(q)$ $E_7(q)$ $E_8(q)$ $F_4(q)$ $G_2(q)$ ${}^3D_4(q^3)$ ${}^2E_6(q^2)$ ${}^2B_2(2^{2n+1})$ ${}^2F_4(2^{2n+1})$ ${}^2G_2(3^{2n+1})$	$E_7(q)$ $E_8(q)$ $F_4(q)$ $G_2(q)$ ${}^3D_4(q^3)$ ${}^2E_6(q^2)$ ${}^2B_2(2^{2n+1})$ ${}^2F_4(2^{2n+1})$ ${}^2G_2(3^{2n+1})$	$E_8(q)$ $F_4(q)$ $G_2(q)$ ${}^3D_4(q^3)$ ${}^2E_6(q^2)$ ${}^2B_2(2^{2n+1})$ ${}^2F_4(2^{2n+1})$ ${}^2G_2(3^{2n+1})$	$F_4(q)$ $G_2(q)$ ${}^3D_4(q^3)$ ${}^2E_6(q^2)$ ${}^2B_2(2^{2n+1})$ ${}^2F_4(2^{2n+1})$ ${}^2G_2(3^{2n+1})$	$G_2(q)$ ${}^3D_4(q^3)$ ${}^2E_6(q^2)$ ${}^2B_2(2^{2n+1})$ ${}^2F_4(2^{2n+1})$ ${}^2G_2(3^{2n+1})$	${}^3D_4(q^3)$ ${}^2E_6(q^2)$ ${}^2B_2(2^{2n+1})$ ${}^2F_4(2^{2n+1})$ ${}^2G_2(3^{2n+1})$	${}^2E_6(q^2)$ ${}^2B_2(2^{2n+1})$ ${}^2F_4(2^{2n+1})$ ${}^2G_2(3^{2n+1})$	${}^2B_2(2^{2n+1})$ ${}^2F_4(2^{2n+1})$ ${}^2G_2(3^{2n+1})$	${}^2F_4(2^{2n+1})$ ${}^2G_2(3^{2n+1})$	${}^2G_2(3^{2n+1})$	$B_n(q)$ $C_n(q)$ $D_n(q)$ ${}^2D_n(q^2)$ ${}^2A_n(q^2)$	$C_n(q)$ $D_n(q)$ ${}^2D_n(q^2)$ ${}^2A_n(q^2)$	$D_n(q)$ ${}^2D_n(q^2)$ ${}^2A_n(q^2)$	${}^2A_n(q^2)$	Z_p C_p p			

- Alternating Groups
- Classical Chevalley Groups
- Chevalley Groups
- Classical Steinberg Groups
- Steinberg Groups
- Suzuki Groups
- Ree Groups and Tits Groups*
- Sporadic Groups
- Cyclic Groups

Alternant*
Symbol
Order†

M_{11}	M_{12}	M_{22}	M_{23}	M_{24}	$J(1), f(11)$	$H1$	$H2$	H/M	I_3	I_4	HS	McL	He	Ru
7920	95040	463320	18209760	244823040	375560	608400	59232960	8677519106 87796240	44352000	998128000	600000720	16942634400		

Sz	$O'N$	O_3	O_2	O_1	HN	Ly	Th	$M(23)$	$M(23)$	F_{24}	F_2	F_3	M
443520 897360	96093 924922	899786 666400	4230832 112400	438776 866	91200000	86100000	867470260	443617034600	4089407473 203000000	1208205708480 860721302400	1208205708480 860721302400	1208205708480 860721302400	1208205708480 860721302400

*The tilde group ${}^{\sim}G$ is not a group of the type but is the product of commutators of G .
† is usually given for binary for type ratios.

*Two sporadic groups and families, alternate names in this legend (all the other names by which they are known) are given in parentheses. All such groups are used to indicate non-triviality. All such groups are listed on the table using the form $A_n(q)$, $B_n(q)$, $C_n(q)$, $D_n(q)$.

The Monster, a.k.a., the Friendly Giant

The largest of all the sporadic finite simple groups, it has order

$$2^{46} \cdot 3^{20} \cdot 5^9 \cdot 7^6 \cdot 11^2 \cdot 13^3 \cdot 17 \cdot 19 \cdot 23 \cdot 29 \cdot 31 \cdot 41 \cdot 47 \cdot 59 \cdot 71$$

or

$$808017424794512875886459904961710757005754368000000000$$

which is approximately $8 \cdot 10^{53}$.